

EDITORIAL COMMENT

Magnetic Resonance Identification of the Ventricular Tachycardia Critical Isthmus

Finding the Needle in the Haystack*

Henry R. Halperin, MD, MAP,
Saman Nazarian, MD
Baltimore, Maryland

Significant strides have been made in development of catheter-based therapeutics for cure of malignant ventricular arrhythmias. Notable initial milestones included the first intracardiac recordings made by Hecht (1), and demonstration of catheter-based cardiac stimulation by Furman and Robinson (2). Later, Wellens et al. (3) demonstrated that programmed electrical stimulation could reliably initiate and terminate ventricular tachycardia (VT) circuits in susceptible individuals. These tools allowed electrophysiologists to grow in their understanding of VT mechanisms, critical pathway, and sites of origin, and potential new therapeutics.

See page 184

In 1978, Josephson et al. (4–6) demonstrated that intraoperative mapping of VT circuits can lead to identification and curative resection of the critical isthmus for VT maintenance. Once excisional cure was demonstrated, techniques were developed for nonsurgical destruction of targeted tissue by high-energy shocks (7–9), and later by radiofrequency energy (10–13). Since then, numerous important contributions have been made to optimize techniques for VT ablation. Of utmost importance is the development of electrophysiology criteria for distinguishing the critical isthmus from bystander areas adjacent to the VT circuit (14). The critical isthmus for VT is often in diseased tissue that exhibits slow conduction properties. However, identification of a slow conduction zone does not necessarily imply vulnerability of the patient to, or participation of, that slow zone in a VT circuit. Electrophysiology maneuvers from the slow zone in question are an integral part of the mechanistic

study of VT and the formulation of an approach to eliminate it. Due to the complex nature of the information derived from such studies, 3-dimensional mapping systems have been developed to summarize the electrical and anatomic information gathered during the study (15,16). Such systems simplify the localization of diseased and slow conduction zones via voltage and activation mapping, but the solution to the VT puzzle is only attainable through the appropriate use and understanding of electrophysiology maneuvers.

Cardiac magnetic resonance (CMR) is capable of providing high resolution images of intramyocardial scar through the late gadolinium enhancement (LGE) technique (17–19). Heterogeneous tissues surrounding areas with dense LGE are hypothesized to be the imaging equivalent of slow conduction zones in patients with ischemic cardiomyopathy (20,21). However, although the relationship between inducibility of VT and extent of LGE in ischemic cardiomyopathy has previously been reported (22,23), little data exists regarding direct electrophysiology measures and maneuvers from heterogeneous sites identified by CMR (24).

In this issue of the *Journal*, Perez-David et al. (25) present the results of a study that compares LGE features using a 1.5-T magnetic resonance scanner, in 18 patients with sustained monomorphic VT, to 18 matched control subjects. The investigators chose to use Yan et al.'s (20) method for defining the infarct core and heterogeneous tissue as regions with signal intensity threshold >3 and between 2 and 3 standard deviations higher than remote normal myocardium, respectively. Continuous heterogeneous corridors were then defined as the presence of heterogeneous tissue in consecutive planes surrounded by scar and connected to normal myocardium by at least 1 side. The underlying hypothesis that slow conduction channels during electrophysiology study would correspond to continuous heterogeneous corridors defined by CMR was then tested. The VT and control groups were similar in scar, heterogeneous tissue, and overall myocardial mass. However, patients in the VT group were more likely to exhibit continuous corridors of heterogeneous tissue. To make the images most suitable for comparison to endocardial voltage maps, the investigators then created 3-dimensional color-coded shells displaying the subendocardial signal-intensity distribution. Continuous corridors of diseased but conducting tissue on endocardial voltage mapping corresponded to continuous corridors of heterogeneous tissue identified by LGE. Of 26 total corridors in 17 of 18 patients with VT, 15 corresponded to a critical isthmus for VT. These results are remarkable. Perez-David et al. (25) have demonstrated the biologically plausible but up to now unproven connection between heterogeneous tissue on CMR and slow conduction zones identified by endocardial mapping. Additionally, they have shown that sites that electrophysiology maneuvers identify as critical VT isthmus sites often reside in heterogeneous tissue identified by CMR.

*Editorials published in the *Journal of the American College of Cardiology* reflect the views of the authors and do not necessarily represent the views of *JACC* or the American College of Cardiology.

From Johns Hopkins Hospital, Baltimore, Maryland. Dr. Halperin is funded by grants R01-HL64795 and R01-HL09461, from the National Institutes of Health. Dr. Nazarian is funded by grant K23-HL089333 from the National Institutes of Health.

These findings have important differences with previous studies. In contrast to findings of previous publications (22,23), the VT and control groups of the current study were not different in scar and heterogeneous tissue mass. This inconsistency may be attributable to differences in patient populations and definition of infarct core and heterogeneous tissue on LGE, or the use of inducibility at electrophysiology study as a surrogate of spontaneous VT. Alternatively, the inconsistency may reflect the fact that a marker of risk in a larger population may not offer adequate resolution in an enriched sample of at-risk individuals. If validated in future studies, the association of continuous corridors of tissue with spontaneous VT may improve the arrhythmic risk stratification of patients with ischemic cardiomyopathy. Another inconsistency with previous studies resides in the location of critical VT zones in the current study versus that performed by Desjardens et al. (24). Desjardens et al. (24) found that the majority of critical VT isthmus sites were within the infarct core as defined by LGE, whereas Perez-David et al. (25) found all critical isthmus sites within heterogeneous tissue. This difference may reflect current CMR shortcomings in scar resolution and volume averaging in the infarct core and border zones. Until higher resolution images capable of resolving the intricate architecture of fibrotic tissue and surrounding healthy myocytes are clinically feasible (26), electrophysiologic characterization of CMR surrogates will be limited by the reality that the currently visualized “infarct core” and “heterogeneous tissue” is a mixture of true infarct core, true heterogeneous tissue, and artifacts, particularly those produced by volume averaging of infarct core with healthy tissue.

The investigators have presented data to support further research in visualization of the substrate for VT. Identification of the infarct core and its associated heterogeneous tissue before the electrophysiology study and integration of such information into mapping systems will likely decrease procedure and fluoroscopy duration by focusing our attention on areas with diseased tissue. However, the electrophysiologist must remember that LGE image signal intensity is very sensitive to the assigned inversion time, poor ECG gating in the setting of arrhythmia, artifacts from fat in the atrioventricular groove or epicardium, and artifacts from respiratory motion. Additionally, volume averaging may represent a perfectly sharp but slanted scar border as “heterogeneous tissue” on LGE. Once such CMR limitations are addressed, we will know where on the LGE image to focus our attention. However, as is evident from the outstanding report by Perez-David et al. (25), solving the VT puzzle will still need an electrophysiologist who is adept at maneuvers to distinguish critical sites from bystanders.

Reprint requests and correspondence: Dr. Henry R. Halperin, Professor of Medicine, Radiology, and Biomedical Engineering, Johns Hopkins Hospital, Blalock 524A, 600 North Wolfe Street, Baltimore, Maryland 21287. E-mail: hhalper@jhmi.edu.

REFERENCES

1. Hecht HH. Potential variations of the right auricular and ventricular cavities in man (the human electrogram). *Proc Am Fed Clin Res* 1947;3:91.
2. Furman S, Robinson G. The use of an intracardiac pacemaker in the correction of total heart block. *Surg Forum* 1958;9:245-8.
3. Wellens HJ, Schuilenburg RM, Durrer D. Electrical stimulation of the heart in patients with ventricular tachycardia. *Circulation* 1972;46:216-26.
4. Josephson ME, Horowitz LN, Farshidi A, Kastor JA. Recurrent sustained ventricular tachycardia. 1. Mechanisms. *Circulation* 1978;57:431-40.
5. Josephson ME, Horowitz LN, Farshidi A, Spear JF, Kastor JA, Moore EN. Recurrent sustained ventricular tachycardia. 2. Endocardial mapping. *Circulation* 1978;57:440-7.
6. Josephson ME, Harken AH, Horowitz LN. Endocardial excision: a new surgical technique for the treatment of recurrent ventricular tachycardia. *Circulation* 1979;60:1430-9.
7. Hartzler GO. Electrode catheter ablation of refractory focal ventricular tachycardia. *J Am Coll Cardiol* 1983;2:1107-13.
8. Puech P, Gallay P, Grolleau R, Koliopoulos N. [Treatment by endocavitary fulguration of recurrent ventricular tachycardia caused by right ventricular dysplasia]. *Arch Mal Coeur Vaiss* 1984;77:826-34.
9. Fontaine G, Tonet JL, Frank R, et al. [Emergency treatment of chronic ventricular tachycardia after myocardial infarction by endocavitary fulguration]. *Arch Mal Coeur Vaiss* 1985;78:1037-43.
10. Budde T, Breithardt G, Borggrefe M, Podczeczek A, Langwasser J. [Initial experiences with high-frequency electric ablation of the AV conduction system in the human]. *Z Kardiol* 1987;76:204-10.
11. Lee MA, Morady F, Kadish A, et al. Catheter modification of the atrioventricular junction with radiofrequency energy for control of atrioventricular nodal reentry tachycardia. *Circulation* 1991;83:827-35.
12. Jackman WM, Wang XZ, Friday KJ, et al. Catheter ablation of accessory atrioventricular pathways (Wolff-Parkinson-White syndrome) by radiofrequency current. *N Engl J Med* 1991;324:1605-11.
13. Kuck KH, Schluter M. Single-catheter approach to radiofrequency current ablation of left-sided accessory pathways in patients with Wolff-Parkinson-White syndrome. *Circulation* 1991;84:2366-75.
14. Stevenson WG, Khan H, Sager P, et al. Identification of reentry circuit sites during catheter mapping and radiofrequency ablation of ventricular tachycardia late after myocardial infarction. *Circulation* 1993;88:1647-70.
15. Gepstein L, Hayam G, Ben-Haim SA. A novel method for nonfluoroscopic catheter-based electroanatomical mapping of the heart. In vitro and in vivo accuracy results. *Circulation* 1997;95:1611-22.
16. Wittkampf FH, Wever EF, Derksen R, et al. Localisa: new technique for real-time 3-dimensional localization of regular intracardiac electrodes. *Circulation* 1999;99:1312-7.
17. McNamara MT, Higgins CB, Ehman RL, Revel D, Sievers R, Brasch RC. Acute myocardial ischemia: magnetic resonance contrast enhancement with gadolinium-DTPA. *Radiology* 1984;153:157-63.
18. Simonetti OP, Kim RJ, Fieno DS, et al. An improved MR imaging technique for the visualization of myocardial infarction. *Radiology* 2001;218:215-23.
19. Kim RJ, Fieno DS, Parrish TB, et al. Relationship of MRI delayed contrast enhancement to irreversible injury, infarct age, and contractile function. *Circulation* 1999;100:1992-2002.
20. Yan AT, Shayne AJ, Brown KA, et al. Characterization of the peri-infarct zone by contrast-enhanced cardiac magnetic resonance imaging is a powerful predictor of post-myocardial infarction mortality. *Circulation* 2006;114:32-9.
21. Roes SD, Borleffs CJ, van der Geest RJ, et al. Infarct tissue heterogeneity assessed with contrast-enhanced MRI predicts spontaneous ventricular arrhythmia in patients with ischemic cardiomyopathy and implantable cardioverter-defibrillator. *Circ Cardiovasc Imaging* 2009;2:183-90.
22. Bello D, Fieno DS, Kim RJ, et al. Infarct morphology identifies patients with substrate for sustained ventricular tachycardia. *J Am Coll Cardiol* 2005;45:1104-8.
23. Schmidt A, Azevedo CF, Cheng A, et al. Infarct tissue heterogeneity by magnetic resonance imaging identifies enhanced cardiac arrhythmia

- susceptibility in patients with left ventricular dysfunction. *Circulation* 2007;115:2006-14.
24. Desjardins B, Crawford T, Good E, et al. Infarct architecture and characteristics on delayed enhanced magnetic resonance imaging and electroanatomic mapping in patients with postinfarction ventricular arrhythmia. *Heart Rhythm* 2009;6:644-51.
25. Perez-David E, Arenal Á, Rubio-Guivernau JL, et al. Noninvasive identification of ventricular tachycardia-related conducting channels using contrast-enhanced magnetic resonance imaging in patients with chronic myocardial infarction: comparison of signal intensity scar mapping and endocardial voltage mapping. *J Am Coll Cardiol* 2011;57:184-94.
26. Ashikaga H, Sasano T, Dong J, et al. Magnetic resonance-based anatomical analysis of scar-related ventricular tachycardia: implications for catheter ablation. *Circ Res* 2007;101:939-47.

Key Words: ablation ■ heterogeneous imaging ■ magnetic resonance imaging ■ ventricular tachycardia.